

Tight Bounds on the Redundancy of Huffman Codes

Soheil Mohajer*, Payam Pakzad*, Ali Kakhbod†

February 1, 2008

1 Introduction

Consider a discrete finite source with N symbols, and with the probability distribution $\mathbf{p} := (u_1, u_2, \dots, u_N)$. It is well-known that the Huffman encoding algorithm [1] provides an optimal prefix code for this source. A D -ary Huffman code is usually represented using a D -ary tree $\mathcal{T}$, whose leaves correspond to the source symbols; The D edges emanating from each intermediate node of $\mathcal{T}$ are labeled with the D letters of the alphabet, and the codeword corresponding to a symbol is the string of labels on the path from the root to the corresponding leaf. Huffman's algorithm is a recursive bottom-up construction of $\mathcal{T}$, where at each time the smallest D probabilities are merged into a new unit, and henceforth represented by an intermediate node in the tree. Throughout this paper, unless D is explicitly specified, we talk about the binary Huffman codes.

Denote by $l(u)$ the length of the path from the root to a node u on the Huffman tree $\mathcal{T}$. Then the expected length of the Huffman code is defined as

$$L(\mathcal{T}) := \sum_{i=1}^k u_i l(u_i). \quad (1)$$

Similarly, the entropy is defined as

$$H(\mathcal{T}) := - \sum_{i=1}^k u_i \log_D(u_i). \quad (2)$$

The Huffman encoding is optimal in the sense that no other code for distribution $\mathbf{p}$ can have a smaller expected length than $L(\mathcal{T})$.

The redundancy $R(\mathcal{T})$ of the code is defined as the difference between the average codeword length $L(\mathcal{T})$, and the entropy $H(\mathcal{T})$ of the source. It is easy to show that the redundancy of the Huffman code is always non-negative and never exceed 1. These bounds on $R(\mathcal{T})$ can be improved if partial knowledge about the source distribution is available. Gallager [2], Johnsen [3], Capocelli and Desantis [4] and [5], Manstetten [6] and Capocelli *et al.* [7] improved the upper bound on the redundancy (of binary Huffman codes) in terms of $p_1 := \max_i u_i$, the probability of the most likely source symbol. On the other hand, in [4] and [8] the problem of upper bounding the redundancy have been addressed when $p_N := \min_i u_i$, the probability of the least likely source symbol, is known. Furthermore Capocelli *et al.* [7] obtained upper bounds on $R(\mathcal{T})$ when both extreme probabilities, p_1 and p_N are known. Also in [8] and [9] upper bound as a function of two least likely source symbol probabilities is derived.

*School of Computer and Communication Sciences, Ecole Polytechnique Fédérale de Lausanne, Switzerland. E-mail: {soheil.mohajer, payam.pakzad}@epfl.ch

†Department of Electrical and Computer Engineering, Isfahan University of Technology, Iran. E-mail: ali_kakhbod@ec.iut.ac.ir




Figure 1: Decomposition of a Huffman tree around an intermediate node u .

Johnsen [3] presented a tight lower bound on $R(\mathcal{T})$ in terms of p_1 when $p_1 \geq 0.4$. Subsequently, such lower bounds were presented by Montgomery and Abrahams [10] for all p_1 . Golić and Obradović [11] extended Johnsen's lower bound on the redundancy of a D -ary Huffman code. The lower bound on $R(\mathcal{T})$ when only p_N is known, is considered in [4]. Furthermore, the problem of lower bounding $R(\mathcal{T})$ for a binary code when the two least likely probabilities are known, is discussed in [4] and [9].

Recently Ye and Yeung [12] presented a simple upper bounds on $R(\mathcal{T})$ in terms of the probability of *any* source symbol, as opposed to the case the least or the most likely probability is given. Using a complicated approach they conjectured the tight upper bound on $R(\mathcal{T})$ for a source containing a symbol with a given probability p , without knowledge about its 'rank in the source distribution'. In this paper we prove this conjecture with a simple approach and prove that this upper bound is tight. Similarly, we present a tight lower bound on $R(\mathcal{T})$ for a source that contains a symbol with given probability p . We further describe all possible sets of distribution which achieve this lower bound. We show that simple extensions of our results lead to the lower bound on the redundancy when either p_1 [10] or p_N [4] are known. We also extend our proof to the D -ary Huffman codes and find the tight lower bound on $R(\mathcal{T})$ when probability of any symbols is known.

2 Preliminaries

In this section we present some definitions and results that will be useful in the rest of the paper.

Let $\mathcal{T} = \mathcal{T}(\mathbf{p})$ be a binary Huffman tree for a source with probability distribution $\mathbf{p} = (u_1, u_2, \dots, u_N)$. For the rest of this paper, we identify each node of a Huffman tree by the probability of the corresponding symbol or unit; this is defined as the sum of the probabilities of all the leaf symbols lying in the sub-tree under that node. For each intermediate node u of $\mathcal{T}$, let Δ_u denote the sub-tree of $\mathcal{T}$ under u , and denote by $u^{-1} * \Delta_u$ its normalized version, where the probability of each node is scaled by a factor of $1/u$, so that the scaled leaf probabilities sum to one. Therefore, $u^{-1} * \Delta_u$ itself is a Huffman tree for a source with probabilities which are given on the leaves of $(u^{-1} * \Delta_u)$. Similarly, denote by Λ_u the Huffman tree on a source with probabilities which are the same as $\mathbf{p}$, except that all the leaf probabilities of $\mathcal{T}$ under u are merged as a single symbol with probability u , which appears as a leaf of Λ_u . Then it is easy to see that Λ_u corresponds to the sub-tree of $\mathcal{T}$ appearing 'above' u . See Figure 1 for a schematic diagram of the relationship between Δ_u and Λ_u with the original Huffman tree $\mathcal{T}$.

The following lemma relates the redundancy of a Huffman tree to the redundancies of the sub-trees $(u^{-1} * \Delta_u)$ and Λ_u .

Lemma 1. For any intermediate node u in a Huffman tree $\mathcal{T}$, we have

$$R(\mathcal{T}) = R(\Lambda_u) + u R(u^{-1} * \Delta_u) \quad (3)$$

Proof. The average length of any Huffman code equals the sum of the probabilities on the intermediate nodes in the corresponding tree. Each intermediate node of $\mathcal{T}$ is either an intermediate node in Λ_u or in $(u^{-1} * \Delta_u)$, where the probabilities in the latter tree need to be scaled back by a factor of u . Therefore we have

$$L(\mathcal{T}) = L(\Lambda_u) + u L(u^{-1} * \Delta_u). \quad (4)$$

Similarly, decomposing the leaf nodes of $\mathcal{T}$ in terms of those on Λ_u and on $(u^{-1} * \Delta_u)$, we get

$$\begin{aligned} H(\mathcal{T}) &= - \sum_{x \in \text{leaf}(\mathcal{T})} x \log x \\ &= - \sum_{x \in \text{leaf}(\Lambda_u) \setminus \{u\}} x \log x - \sum_{y \in \text{leaf}(u^{-1} * \Delta_u)} u y \log u y \\ &= - \left(\sum_{x \in \text{leaf}(\Lambda_u)} x \log x - u \log u \right) - \left(u \sum_{y \in \text{leaf}(u^{-1} * \Delta_u)} y \log y + u \log u \right) \\ &= H(\Lambda_u) + u H(u^{-1} * \Delta_u) \end{aligned} \quad (5)$$

and the desired result will be obtained immediately from (4) and (5). $\blacksquare$

3 Upper Bound

In this section we provide a tight upper bound for the redundancy of the Huffman code for a source containing a symbol with a given probability p . Note that p need not be the probability of the most or the least likely symbol.

Theorem 1 (Tight Upper Bound on Huffman Redundancy). Consider the Huffman code for a source with finite alphabet, which includes a symbol with probability p , but is otherwise arbitrary. The redundancy of this code is upper bounded by

$$R_{\max}(p) := \begin{cases} 2 - p - \mathcal{H}(p), & \text{if } 0.5 \leq p < 1 \\ 1 + p - \mathcal{H}(p), & \text{if } 0 \leq p < 0.5 \end{cases} \quad (6)$$

where $\mathcal{H}(p) := -p \log p - (1-p) \log(1-p)$ is the binary entropy function. Furthermore, this bound is tight, so there are sequences of sources whose Huffman redundancies converge to $R_{\max}(p)$.

Before we prove this theorem, we shall review some previous related results, which will be used in our proof.

Our result improves the following bound obtained in [12], and in fact proves a conjecture for the tightest upper bound given in the same paper:

Theorem 2. Let p be the probability of any source symbol. Then the redundancy of the corresponding Huffman code is upper bounded by

$$R_{\text{upperbound}}(p) := \begin{cases} 2 - p - \mathcal{H}(p), & \text{if } 0.5 \leq p < 1 \\ 0.5, & \text{if } \pi_0 \leq p < 0.5 \\ 1 + p - \mathcal{H}(p), & \text{if } p < \pi_0 \end{cases} \quad (7)$$

where $\pi_0 \simeq 0.18$ is the smallest root of equation $1 + p - \mathcal{H}(p) = 0.5$.




Figure 2: Upper bounds of Theorems 1, 2 and 3, on the redundancy of a source containing a symbol with probability p .

We will skip the proof here and refer the reader to the original paper [12]. This upper bound is tight when $p \geq 0.5$ or $p < \pi_0 \simeq 0.18$, but as also suggested in [12], it is not tight for the central region $0.18 < p < 0.5$. Thus, in our proof of Theorem 1 we will only consider this central region and obtain a tight bound for the redundancy.

We will also use the following upper bound on the redundancy of a source whose *most likely* symbol probability is known. A more precise form of this bound is included in [6], and we refer the interested readers to that work for details and proof.

Theorem 3. Let p_1 be the probability of the most likely symbol of a source. Then the Huffman redundancy of this source is upper bounded by the following function:

$$f(p_1) = \begin{cases} 2 - p_1 - \mathcal{H}(p_1) & \text{if } 0.5 \leq p_1 < 1 \\ 3 - 5p_1 - \mathcal{H}(2p_1) & \text{if } \pi_1 \leq p_1 < 0.5 \\ \gamma & \text{if } p_1 \leq \pi_1 \end{cases} \quad (8)$$

where $\gamma = R_{\max}(\frac{1}{3}) = 1 + 1/3 - \mathcal{H}(1/3) \simeq 0.415$, and $\pi_1 \simeq 0.491$ is a root of $3 - 5p_1 - \mathcal{H}(2p_1) = \gamma$, see Figure 2.

Finally we will need the following lemma in our proof of Theorem 1.

Lemma 2. Let p_1 be the probability of the most likely letter in a source, which also contains another letter with probability q . If $p_1 \geq \pi_1 \simeq 0.491$ and $q \geq \pi_0 \simeq 0.18$, then $l(p_1) = 1$, i.e. the length of the codeword corresponding to the most likely symbol is 1.

Proof. We first note that as long as $p_1 > \frac{1}{3}$, the length $l(p_1)$ cannot be larger than 2; otherwise, there would be at least two independent intermediate nodes x and y on the Huffman tree with lengths less than p_1 . Therefore x and y both would have probabilities at least as large as p_1 ; but this is a contradiction since $p_1 + x + y \geq 3p_1$ cannot exceed 1.




Figure 3: A Huffman tree with $l(p_1) = 2$.

Suppose next that $l(p_1) = 2$. Then there can be no codeword of length 1 in the code, since p_1 is the largest probability. Therefore, the corresponding tree has a structure as in Figure 3.

Note first that $v_1 + v_2 \geq p_1$ since $l((v_1 + v_2)) = 1 < l(p_1) = 2$. We will next show that $u \geq q$. Clearly, if q lies in the sub-tree under u , the assertion follows. Suppose then that q lies in the sub-tree under v_1 . Note next that, by the construction of the Huffman tree, either $p_1, u \geq v_1, v_2$, or $p_1, u \leq v_1, v_2$. The second alternative cannot happen, since then $1 \geq p_1 + v_1 + v_2 \geq 3p_1 \geq 3\pi_1 \simeq 1.47$ would be a contradiction. Therefore $u \geq v_1 \geq q$.

Combining these relationships we get

$$\pi_1 \leq p_1 \leq v_1 + v_2 = 1 - p_1 - u \leq 1 - p_1 - q \leq 1 - \pi_1 - \pi_0,$$

which is a contradiction, since $\pi_1 \simeq 0.491$ and $1 - \pi_0 - \pi_1 \simeq 0.329$. $\blacksquare$

Proof of Theorem 1. As stated before, when $p \leq \pi_0 \simeq 0.18$ or $p \geq 0.5$ our bound coincides with the bound of Theorem 2. It remains to show that for $\pi_0 < p < 0.5$, the redundancy of a Huffman code for a source which contains a symbol with probability p cannot be larger than $R_{\max}(p)$.

We prove this claim using an argument on p_1 , the probability of the most likely symbol in $\mathbf{p}$. First note that, if $p = p_1$ is the most likely symbol, then from Theorem 3, $R(\mathcal{T}) \leq f(p) \leq R_{\max}(p)$. Suppose then that $p < p_1$. Clearly $p_1 \leq 1 - p$ since p and p_1 are probabilities in the same distribution $\mathbf{p}$. Once again, from Theorem 3, if $p_1 \leq \pi_1$, then the Huffman redundancy of $\mathbf{p}$ cannot exceed $R_{\max}(p)$ for any p , since $R_{\max}(p) \geq R_{\max}(\frac{1}{3}) \geq f(p)$ for all $p \leq \pi_1$.

Suppose then that $p_1 > \pi_1$. Let $\mathcal{T}$ be a Huffman tree for $\mathbf{p}$, which contains $p > \pi_0$ as a leaf. Then by Lemma 2 we have $l(p_1) = 1$. Thus p should appear in the sub-tree under the intermediate node of probability $1 - p_1$, i.e. $p \in \Delta_{1-p_1}$ and hence $q := \frac{p}{1-p_1} \in (1 - p_1)^{-1} * \Delta_{1-p_1}$. Then by Lemma 1, we have

$$R(\mathcal{T}) = 1 - \mathcal{H}(p_1) + (1 - p_1)R((1 - p_1)^{-1} * \Delta_{(1-p_1)}) \quad (9)$$

We then upper bound the term $R((1 - p_1)^{-1} * \Delta_{(1-p_1)})$ using Theorem 2. Note that $\pi_0 < q \leq 1$, since p was assumed to be greater than π_0 . We consider the following two cases on the possible values of $q = \frac{p}{1-p_1}$:

- Case 1: $\pi_0 < q \leq 0.5$. From Theorem 2, $R((1 - p_1)^{-1} * \Delta_{(1-p_1)}) \leq \frac{1}{2}$. Then using (9) we get

$$R(\mathcal{T}) \leq 1 - \mathcal{H}(p_1) + \frac{(1 - p_1)}{2} \quad (10)$$

The right-hand-side of the above inequality is a convex function of $p_1 \in (\pi_1, 1 - \pi_0)$ and it is easy to verify that it takes its maximum value at the boundary point $p_1 = 1 - \pi_0 \simeq 0.82$. Then we have

$$\begin{aligned}
R(\mathcal{T}) &\leq \max_{p_1 \in (\pi_1, 1 - \pi_0)} \left(1 - \mathcal{H}(p_1) + \frac{(1 - p_1)}{2} \right) \\
&= \left(1 - \mathcal{H}(\pi_0) + \frac{\pi_0}{2} \right) \simeq 0.410 \\
&< \min_{p \in (\pi_0, 0.5)} R_{\max}(p) \\
&= R_{\max}\left(\frac{1}{3}\right) \simeq 0.415
\end{aligned}$$

- Case 2: $0.5 < q \leq 1$. Define $\beta := 1 - q$, so that $\mathcal{H}(\beta) = \mathcal{H}(q)$. Then once again, using the upper bound of Theorem 2 in (9) we get

$$\begin{aligned}
R(\mathcal{T}) &\leq 1 - \mathcal{H}(p_1) + (1 - p_1)R_{\text{upperbound}}(q) \\
&= 1 - \mathcal{H}(p_1) + (1 - p_1)(2 - q - \mathcal{H}(q)) \\
&\leq 1 - \mathcal{H}(p) + (1 - p_1)(2 - q - \mathcal{H}(\beta)) \\
&= 1 - \mathcal{H}(p) + p + (1 - p_1)(2 - 2q - \mathcal{H}(q)) \\
&= R_{\max}(p) - (1 - p_1)(\mathcal{H}(\beta) - 2\beta) \\
&\leq R_{\max}(p)
\end{aligned}$$

where in the third line we have used the fact that $\mathcal{H}(p_1) \geq \mathcal{H}(p)$ since $p \leq p_1 \leq 1 - p$, and the last inequality follows from the fact that $(\mathcal{H}(\beta) - 2\beta) \geq 0$ for $0 \leq \beta < 0.5$.

Thus we have shown that $R(\mathcal{T})$ is upper bounded by $R_{\max}(p)$ for all p . It only remains to show the tightness of the bound. It is easy to check that the redundancy of a source with distribution $\mathbf{p}_\epsilon := ((1 - \epsilon)(1 - p), p, \epsilon(1 - p))$ is

$$R(\mathbf{p}_\epsilon) = 1 + p - \mathcal{H}(p) - (1 - p)(\mathcal{H}(\epsilon) - \epsilon), \quad (11)$$

which tends to $R_{\max}(p) = 1 + p - \mathcal{H}(p)$ as ϵ goes to zero. This completes the proof. $\blacksquare$

4 Lower Bound

In this section we provide a lower bound for the redundancy of the Huffman code for a source that contains a symbol with a given probability p . As we will see, the redundancy can be zero only if p is dyadic, i.e $p = 2^{-l}$ for some integer l . We will also show that our bound is tight and describe all possible sets of distributions which achieve this redundancy.

Theorem 4 (Tight Lower Bound on Huffman Redundancy). Consider the Huffman code for a source with finite alphabet, which includes a symbol with probability p , but is otherwise arbitrary. The redundancy of this code is lower bounded by

$$R_{\min}(p) := mp - \mathcal{H}(p) - (1 - p) \log(1 - 2^{-m}), \quad (12)$$

where $m > 0$ takes either of the values $\lfloor -\log p \rfloor$ or $\lceil -\log p \rceil$ which minimizes the expression. Furthermore, this bound is tight, i.e. there exist sources containing a symbol with probability p , whose Huffman redundancies equal $R_{\min}(p)$.




Figure 4: Canonical structure for minimum-redundancy Huffman trees that contain a symbol with probability p .

Proof. We first note that, for the purposes of minimizing the redundancy, it suffices to only consider a simple class of probability distributions, which is depicted in Figure 4. To see this, let u be any intermediate node in a Huffman tree $\mathcal{T}$, which does not contain p in the sub-tree under it, i.e. $p \notin \Delta_u$. Then from Lemma 1,

$$R(\mathcal{T}) = R(\Lambda_u) + uR(u^{-1} * \Delta_u) \geq R(\Lambda_u). \quad (13)$$

Therefore Λ_u is a Huffman tree containing a leaf with probability p , whose redundancy is at most equal to that of $\mathcal{T}$. This argument can be repeated until $\mathcal{T}$ is converted into the form of Figure 4.

Suppose then, without loss of generality, that $\mathcal{T}$ has this simplified form. Now let $x_i = \alpha_i \cdot (1 - p)$, where $\sum_{i=1}^m \alpha_i = 1$. We can write the following expressions for the expected length of the code and the entropy of the source:

$$\begin{aligned} L(\mathcal{T}) &= \sum_{i=1}^m x_i \cdot i + pm \\ &= 1 + (m-1)p + (1-p) \sum_{i=1}^m (i-1)\alpha_i \end{aligned}$$

and

$$\begin{aligned} H(\mathcal{T}) &= - \sum_{i=1}^m (1-p)\alpha_i \log((1-p)\alpha_i) - p \log p \\ &= \mathcal{H}(p) + (1-p)H(\alpha_1, \alpha_2, \dots, \alpha_m). \end{aligned}$$

Thus,

$$\begin{aligned} R(\mathcal{T}) &= 1 + (m-1)p - \mathcal{H}(p) + (1-p) \left[\sum_{i=1}^m (i-1)\alpha_i - H(\alpha_1, \alpha_2, \dots, \alpha_m) \right] \\ &= 1 + (m-1)p - \mathcal{H}(p) + (1-p)f(\alpha_1, \alpha_2, \dots, \alpha_m) \end{aligned} \quad (14)$$

where $f(\alpha_1, \alpha_2, \dots, \alpha_m) := \sum_{i=1}^m (i-1)\alpha_i - H(\alpha_1, \alpha_2, \dots, \alpha_m)$. For each fixed length m of the tree, we wish to find the values of α_i which minimize $R(\mathcal{T})$. Note first that the minimizing probability vector $(\alpha_1, \dots, \alpha_m)$ must be an interior point in the probability simplex, since in the case $\alpha_m = 0$, we can remove $x_m = 0$ from the distribution, —replacing m with $(m-1)$,— and lower the redundancy. Next note that $R(\mathcal{T})$ depends on the α_i 's only through $f(\cdot)$. Writing $\alpha_m =$




Figure 5: Lower bound and Upper bound on the redundancy of a source containing a symbol with probability p .

$1 - \sum_{i=1}^{m-1} \alpha_i$, and differentiating f with respect to $\alpha_1, \dots, \alpha_{m-1}$ we get

$$\begin{aligned} \frac{\partial f}{\partial \alpha_i} &= (i-1-m) + 1 + \log \alpha_i - 1 - \log(1 - \alpha_1 - \dots - \alpha_{m-1}) \\ &= (i-1) + \log \left(\frac{\alpha_i}{1 - \alpha_1 - \dots - \alpha_{m-1}} \right) \end{aligned}$$

Setting $\frac{\partial f}{\partial \alpha_i} = 0$ for $i = 1, \dots, m-1$ we get

$$\frac{\alpha_i}{1 - \alpha_1 - \dots - \alpha_{m-1}} = 2^{m-i}.$$

The unique solution of the above system of equations is

$$\alpha_i = \frac{2^{m-i}}{2^m - 1} \quad (15)$$

Plugging these values into (14) and after straightforward manipulations, we get

$$R(T) = mp - \mathcal{H}(p) - (1-p) \log(1 - 2^{-m}).$$

This is readily seen to be a convex function of m . To minimize, we differentiate with respect to m and set the derivative equal to zero.

$$\frac{\partial}{\partial m} R(T) = p - \frac{1-p}{2^m - 1} = 0,$$

yielding $m = -\log p$. Since m needs to be an integer and by convexity, one of the two neighboring integers $\lfloor -\log p \rfloor$ or $\lceil -\log p \rceil$ will give the minimum. It remains to verify that the α_i values of (15) are consistent with a Huffman tree of the form in Figure 4. A necessary and sufficient condition for this is that $p \leq x_{m-1} = (1-p)\alpha_{m-1}$. It is then easy to see that the chosen value of m in $\{\lfloor -\log p \rfloor, \lceil -\log p \rceil\}$ which minimizes (12), results in an α_{m-1} coefficient which satisfies this condition. ■

Remark 1. An alternative way to describe the minimizing value of m in Theorem 4 is as follows: The minimizing m satisfies

$$\beta_{m-1} \leq p \leq \beta_m \quad (16)$$

where $\beta_0 := 1$ and β_k is given by

$$\beta_k := \left(1 + 1/\log \left(1 + \frac{1}{2^{k+1} - 2} \right) \right)^{-1}.$$

This equation is obtained by equating the values of $R(\mathcal{T})$ from (12) for two consecutive integers. It is easy to see that β_k is a descending sequence, converging to 0 as k grows to infinity, so that for any $p \in (0, 1)$ there exists a unique m satisfying (16). The first few β_k 's are $\beta_1 = 0.369, \beta_2 = 0.182, \beta_3 = 0.091, \dots$ and are displayed in Figure 5.

Remark 2. From (12), the lower bound $R_{\min}(p)$ can be zero if and only if $p = 2^{-m}$ is dyadic. In that case, from (15) $x_i = (1 - p)\alpha_i = 2^{-i}$. Therefore the entire distribution is dyadic.

Remark 3. The proof of Theorem 4 essentially describes all the source distributions that contain a symbol of probability p , and achieve the lower bound $R_{\min}(p)$ on the redundancy. As argued, the Huffman trees for all such distributions have a ‘backbone’ of form of Figure 4, with probabilities that are uniquely determined by the theorem. Any such tree which extends beyond this unique backbone must satisfy the inequality of (13) with equality, i.e. $R(x_i^{-1} * \Delta_{x_i})$ must be zero for all intermediate x_i 's. From Remark 2 above, this can happen only if the corresponding distributions for the sub-trees $x_i^{-1} * \Delta_{x_i}$ are dyadic. Thus, all the distributions containing a symbols of probability p , which achieve the lower bound $R_{\min}(p)$ can be obtained in the following way:

Start with the backbone distribution described in Theorem 4. At any time, choose a leaf node other than p and split its probability in half. Each tree during this process is a Huffman tree with redundancy $R_{\min}(p)$.

In the remainder of this section, we extend the results of Theorem 4 to the cases when the given probability p corresponds to the most, or the least likely symbol.

Suppose first that p is constrained to be the probability of the most likely symbol of a source. Let $\mathbf{p}$ be a distribution as prescribed by Theorem 4, which achieves the minimum redundancy $R_{\min}(p)$, but in which p is not necessarily the maximum probability. Then, by the argument of the Remark 3 above, each symbol probability of $\mathbf{p}$ other than p can be successively split into two halves. We can repeat this process until p becomes the largest value in the distribution. Thus we have the following result:

Theorem 5. (*c.f. Theorem 2 in [10]*) A tight lower bound for the Huffman redundancy of a source whose maximum symbol probability is p_1 is $R_{\min}(p_1)$, as defined in Theorem 4.

Next suppose that $p \leq 0.5$ is constrained to be the probability of the least likely symbol of a source. The argument used in the proof of Theorem 4 can be extended to this case, since even without this additional constraint, we found that in order to achieve the minimum redundancy, p should have the maximum length in the canonical case of Figure 4. We only need to apply the more stringent constraint

$$p \leq x_m = (1 - p)/(2^m - 1). \quad (17)$$

From the convexity of the function $R(\mathcal{T})$ in (14), the minimum will be achieved at one of the points on the boundary of the constraint set which are closest




Figure 6: Lower bound for the Huffman redundancy of a source containing a symbol with probability p .

to the optimal m . Of the two neighboring integers to the optimal m , only $m = \lfloor \log \frac{1}{p} \rfloor$ satisfies the constraint of (17). On the other hand, satisfying the constraint (17) with equality corresponds to a Huffman tree in form of Figure 4, in which $x_m = p$. But, as argued in the remarks above, the redundancy of any such code is lower bounded by $R_{\min}(2p)$, since x_m and p can be merged together with no loss in redundancy, and the resulting tree has a leaf with probability $2p$. Therefore the smallest achievable redundancy is the minimum of $R_{\min}(2p)$ and the value of (14) for $m = \lfloor \log \frac{1}{p} \rfloor$. Finally, it is easy to see that

$$p \lfloor \log \frac{1}{p} \rfloor - \mathcal{H}(p) - (1-p) \log \left(1 - 2^{-\lfloor \log \frac{1}{p} \rfloor} \right) \leq \\ 2p \lfloor \log \frac{1}{2p} \rfloor - \mathcal{H}(2p) - (1-2p) \log \left(1 - 2^{-\lfloor \log \frac{1}{2p} \rfloor} \right).$$

Thus we have proven the following:

Theorem 6. (*c.f. Theorem 2 in [4]*) A tight lower bound for the Huffman redundancy of a source whose maximum symbol probability is p_N , is the smaller of the following two functions:

$$p \lfloor \log \frac{1}{p} \rfloor - \mathcal{H}(p) - (1-p) \log \left(1 - 2^{-\lfloor \log \frac{1}{p} \rfloor} \right), \quad \text{and} \quad (18)$$

$$2p \lfloor \log \frac{1}{2p} \rfloor - \mathcal{H}(2p) - (1-2p) \log \left(1 - 2^{-\lfloor \log \frac{1}{2p} \rfloor} \right). \quad (19)$$

Figure 6 plots the lower bounds of Theorems 4 and 6 as a function of the fixed probability p .

4.1 Extension to the D -ary Huffman Codes

The method of Section 4 can be extended to obtain tight lower bounds for the D -ary Huffman codes. By using the same argument as in the binary case, we can show that the redundancy of a Huffman tree decreases by merging the




Figure 7: Canonical structure for minimum-redundancy D -ary Huffman trees that contain a symbol with probability p .

symbols in the sub-tree under an intermediate node. Thus, for the purposes of minimizing the redundancy, it suffices to only consider Huffman trees with a structure as in Figure 7.

The following lemma shows further that in the minimum-redundancy structure of Figure 7, all the probabilities in the same level must be equal.

Lemma 3. In any minimum-redundancy D -ary Huffman tree of given in Figure 7, we must have $x_{i,l} = x_{j,l}$ for all $l = 1, \dots, m$, and for all $i, j = 2, \dots, D$.

Proof. If $x_{i,l} \neq x_{j,l}$ for some i, j and l , then we replace all the probabilities in that level by their average $\bar{x}_l := \frac{1}{D-1} \sum_k x_{k,l}$, and show that the new tree is a Huffman tree with strictly smaller redundancy. Note first that since the average is between the maximum and minimum probabilities in that level, the new probabilities are still consistent with the same Huffman structure, i.e. $x_{i,l+1} \leq \min_k x_{k,l} < \bar{x}_l < \max_k x_{k,l} \leq x_{j,l-1}$. Next note that the average length of the tree remains fixed after this process. The entropy of the original tree can be written as

$$H(\mathcal{T}) = h_0 + (D-1)\bar{x}_l H_D\left(\frac{x_{2,l}}{(D-1)\bar{x}_l}, \dots, \frac{x_{D,l}}{(D-1)\bar{x}_l}\right),$$

where h_0 is the contribution from the probabilities in the other levels, and $H_D(\cdot)$ is the base- D entropy function, which is uniquely maximized with the uniform distribution. Therefore the entropy is maximized, —and the redundancy minimized,— with the proposed replacement $x_{i,l} \leftarrow \bar{x}_l$. ■

Using Lemma 3 and an argument identical to the one in the proof of Theorem 4 we get the following result:

Theorem 7. The redundancy of a D -ary Huffman code containing a letter with probability p is tightly lower bounded by

$$R_{\min,D} = mp - \mathcal{H}_D(p) - (1-p) \log_D(1 - D^{-m}) \quad (20)$$

where m is either $\lfloor -\log_D p \rfloor$ or $\lceil -\log_D p \rceil$ which minimizes the above expression, and $\mathcal{H}_D(p) := \mathcal{H}(p)/\log(D)$ is the D -ary entropy function.

5 Conclusion

In this paper we have obtained tight upper and lower bounds on the redundancy of the Huffman code for a source, for which the probability of one of the symbols

is known. Our upper bound proves a conjecture of [12], and our lower bound extends and completes several earlier partial results. We have further discussed the explicit form of the distributions that achieve each of these bounds. Our arguments can be extended to the case of the D -ary Huffman codes, and some of these extensions are included in this paper.

References

- [1] D. A. Huffman, “A Method for the construction of minimum-redundancy codes,” *Proc. IRE*, vol. 40, no. 2, pp. 1098–1101, Sept. 1952.
- [2] R. G. Gallager, “Variations on a theme by Huffman,” *IEEE Trans. Inform. Theory*, vol. 24, no. 6, pp. 668–674, Nov. 1978.
- [3] O. Johnsen, “On the redundancy of binary Huffman codes,” *IEEE Trans. Inform. Theory*, vol. 26, no. 2, pp. 220–222, Mar. 1980.
- [4] R. M. Capocelli and A. D. Santis, “New bounds on the redundancy of Huffman codes,” *IEEE Trans. Inform. Theory*, vol. 37, no. 4, pp. 1095–1104, July 1991.
- [5] —, “Tight upper bounds on the redundancy of Huffman codes,” *IEEE Trans. Inform. Theory*, vol. 35, no. 5, pp. 1084–1091, Sept. 1989.
- [6] Manstetten, “Tight bounds on the redundancy of Huffman codes,” *IEEE Trans. Inform. Theory*, vol. 38, no. 1, pp. 144–151, Jan. 1992.
- [7] R. G. R. M. Capocelli and I. J. Taneja, “Bounds on the redundancy of Huffman codes,” *IEEE Trans. Inform. Theory*, vol. 32, no. 6, pp. 854–857, Nov. 1986.
- [8] R. D. Prisco and A. D. Santis, “On the redundancy achieved by Huffman codes,” *Inform. Sci.*, vol. 88, pp. 131–148, Jan. 1996.
- [9] R. W. Yeung, “Local redundancy and progressive bounds on the redundancy of a Huffman code,” *IEEE Trans. Inform. Theory*, vol. 37, no. 3, pp. 687–691, May 1991.
- [10] B. Montgomery and J. Abrahams, “On the redundancy of optimal binary prefix-condition codes for finite and infinite sources,” *IEEE Trans. Inform. Theory*, vol. 33, no. 1, pp. 156–160, Jan. 1987.
- [11] J. Golić and M. Obradović, “A lower bound on the redundancy of D -ary Huffman codes,” *IEEE Trans. Inform. Theory*, vol. 33, no. 6, pp. 910–911, Nov. 1987.
- [12] C. Ye and R. W. Yeung, “A simple upper bound on the redundancy of Huffman codes,” *IEEE Trans. Inform. Theory*, vol. 48, no. 7, pp. 2132–2138, July 2002.